

VOAs generated by two conformal vectors whose τ -involutions generate S_3 .

Masahiko Miyamoto *

Institute of Mathematics
University of Tsukuba
Tsukuba 305, Japan

Dedicated to Professor Koichiro Harada for his 60th birthday.

Abstract

We determined the inner products of two conformal vectors with central charge $\frac{1}{2}$ whose τ -involutions generate S_3 if none of τ -involutions are trivial. We also see that a subVA generated by such conformal vectors is a VOA with central charge $\frac{1}{2} + \frac{20}{21}$ or $\frac{4}{5} + \frac{6}{7}$ and has a Griess algebra of dimension three or four, respectively.

1 Introduction

A vertex operator algebra (shortly VOA) is a mathematical object for a 2-dimensional conformal field theory, but it comes from the Moonshine conjecture to explain mysterious properties of Monster simple group $\mathbb{M}$ [CN]. As an answer to this conjecture, Frenkel, Lepowsky and Meurman construct the moonshine VOA $V^\natural = \bigoplus_{i=0}^{\infty} V_i^\natural$ whose full automorphism group is the Monster simple group $\mathbb{M}$, [FLM]. Its weight two subspace $V_2^\natural$ coincides with a commutative (non-associative) algebra (called the monstrous Griess algebra) of dimension 196884 constructed by Griess in order to construct the Monster simple group [Gr]. This algebra is studied from a group theoretic point of view. One of the important results is that each $2A$ -involution θ defines a unique idempotent e_θ (called an axis) of the monstrous Griess algebra such that the inner product $\langle e_\theta, e_\phi \rangle$ is uniquely determined by the conjugacy classes of $\theta\phi$, see [Co].

From the view point of vertex operator algebras, the author showed that an involutive automorphism τ_e comes arise from a rational conformal vector with central charge $\frac{1}{2}$ in a

*Supported by the Grants-in-Aids for Scientific Research, No. 13440002 and No. 12874001, The Ministry of Education, Science and Culture, Japan.

Griess algebra, namely, if e generates a rational VOA $L(\frac{1}{2}, 0)$ called an Ising model. then one can define an involutive automorphism τ_e of V by

$$\tau_e : \begin{cases} 1 & \text{on } W_0 \oplus W_{\frac{1}{2}} \\ -1 & \text{on } W_{\frac{1}{16}}, \end{cases}$$

where W_h denotes the sum of all irreducible $\text{VA}(e)$ -modules isomorphic to $L(\frac{1}{2}, h)$ and $\text{VA}(e)$ is a subVOA generated by e .

In the monstrous Griess algebra, it is easy to check that such a conformal vector e (with central charge $\frac{1}{2}$) is corresponding to an axis and τ_e is a $2A$ -involution. Even in a (general) Griess algebra, the inner products are very important and the author showed that if two involutions τ_e and τ_f commute, that is, $\tau_e\tau_f$ is of order two, then the inner product $\langle e, f \rangle$ is 0 or $\frac{1}{32}$. In the moonshine VOA $V^\natural$, $\tau_e\tau_f$ is a $2A$ -involution if and only if $\langle e, f \rangle = \frac{1}{32}$ and a $2B$ -involution if and only if $\langle e, f \rangle = 0$. Conversely, if $\langle e, f \rangle = 0$, then τ_e and τ_f commute. One of the mysteries about the Monster simple group is a property of $2A$ -involutions. The $2A$ -involutions satisfy several interesting properties. For example, they satisfy a 6-transposition property, that is, $|\tau_e\tau_f| \leq 6$ for any $2A$ -involutions τ_e and τ_f . There is also a mysterious relation with E_8 Dynkin diagram, which was observed by McKay. Namely, the conjugacy classes of $\tau_e\tau_f$ for distinct $2A$ -involutions τ_e and τ_f are

$$2A, 3A, 4A, 5A, 6A, 3C, 4B \text{ and } 2B$$

and these numbers are multiplicities of roots in a maximal root in E_8 -root system. Another interesting topic is a $Y_{5,5,5}$ -diagram or the set of 26 involutions whose graph coincides with the incident graph given by 13 lines and 13 points of the projective plane of order three and the Bimonster $\mathbb{M} \wr \mathbb{Z}_2$ contains such a set as generators, [At]. If we restrict them to the Monster simple group $\mathbb{M}$, which contains 21 $2A$ -involutions whose graph is the incident graph of 12 lines and 9 points in the Affine plane of order three, where a vertex is a $2A$ -involution τ_e and an edge $\tau_e - \tau_f$ means $|\tau_e\tau_f| = 3$ and no edge between τ_e and τ_f implies that $\tau_e\tau_f$ is of order two, see [Mi1]. Since the author has already studied the case without edge in [Mi2], it becomes very important to treat the case with an edge, that is, the relation between two conformal vectors e and f such that $\tau_e\tau_f$ is of order three.

In this paper, we will study such a pair (e, f) in a (general) Griess algebra with a positive definite invariant bilinear form $\langle \cdot, \cdot \rangle$. We also add one more assumption that none of τ -involutions are trivial. We should note that if $\tau_e = 1$, then we can define an automorphism σ_e of different type by

$$\tau_e : \begin{cases} 1 & \text{on } W_0 \\ -1 & \text{on } W_{\frac{1}{2}} \end{cases}$$

and $\langle \sigma_e | e \in \text{Con}(\frac{1}{2}), \tau_e = 1 \rangle$ is a normal 3-transposition subgroup of the full automorphism group of V and such 3-transposition groups are studied by [F], [GH] and [KMi], where $\text{Con}(\frac{1}{2})$ is the set of all conformal vector with central charge $\frac{1}{2}$. Moreover, if σ_e is

also trivial, then V is a tensor product of a sub VOA $\text{VA}(e)$ and some subVOA W . Since we are interested in the property of group generated by τ -involutions, we get rid of these cases and assume that $\tau_e \neq 1$ for any conformal vector e .

In the Monster group, if a product of two $2A$ -involutions has order three, then it is a $3A$ -trality or a $3C$ -trality. For such conformal vectors in the monstrous Griess algebra $V_2^{\natural}$, the inner products are $\frac{13}{2^{10}}$ and $\frac{1}{2^8}$ corresponding to $3A$ -trality or $3C$ -trality, respectively, see [Co]. We will show that this result is true for general VOAs if none of τ -involutions are trivial, that is, there are only two possibilities of inner products $\langle e, f \rangle$. Moreover, we will study a subVA $\text{VA}(e, f)$ generated by e and f . For example, we will show that $\text{VA}(e, f)$ is a vertex operator algebra. We note that the notation $\text{VA}(e, f)$ denotes a vertex subalgebra generated by e and f and we don't assume the existence of Virasoro element. Since each edge in the graph of 21 involutions is corresponding to a $3A$ -trality, we focus our attention on the case where the inner product is $\frac{13}{2^{10}}$. In this case, $\text{VA}(e, f)$ is a VOA with central charge $\frac{58}{35}$ and $\dim(\text{VA}(e, f))_2 = 4$. Moreover, $\text{VA}(e, f)$ contains $L(\frac{1}{2}, 0) \otimes L(\frac{81}{70}, 0)$ and also $L(\frac{4}{5}, 0) \otimes L(\frac{6}{7}, 0)$, where $L(c, 0)$ denotes a simple Virasoro VOA with central charge c and $L(c, h)$ denotes its irreducible module with highest weight h . In particular, a Griess algebra of $\text{VA}(e, f)$ coincides with the subalgebra generated of e and f as a subalgebra of Griess algebra. We will call this a VOA of τ -involution type A_2 . We may view $L(\frac{1}{2}, 0)$ as a VOA of involution type A_1 .

Acknowledgment

The author wishes to thank R. L. Griess for his helpful advices.

2 Setting and products in Griess algebra

This paper is a continuation of [Mi2] and we will adopt the notation from it. Since our interest is a finite automorphism group, we will treat a simple VOA $(V, Y, \mathbf{1}, \omega)$ over the real number field $\mathbb{R}$ and $\mathbb{C}V$ denotes its complexification $\mathbb{C} \otimes_{\mathbb{R}} V$. For $v \in V$, $Y(v, z) = \sum_{n \in \mathbb{Z}} v(n)z^{-n-1}$ denotes the vertex operator of v .

Assume the following conditions:

- (1) $V = \bigoplus_{n=0}^{\infty} V_n$, $V_0 = \mathbb{R}\mathbf{1}$ and $V_1 = 0$.

A VOA of this type is called OZ(one zero)-type. Since $\dim V_0 = 1$ and $V_1 = 0$, there is a unique invariant bilinear form $\langle \cdot, \cdot \rangle$ on V satisfying $\langle \mathbf{1}, \mathbf{1} \rangle = 1$. We also assume

- (2) $\langle \cdot, \cdot \rangle$ is definite on V_n for each n .

In particular, V_2 has a positive definite bilinear form $\langle \cdot, \cdot \rangle$ satisfying $\langle a(1)b, c \rangle = \langle b, a(1)c \rangle$ for $a, b, c \in V_2$, where $\langle a, b \rangle$ is given by $\langle a, b \rangle \mathbf{1} = a(3)b \in \mathbb{R}\mathbf{1}$. A VOA of this type is called a moonshine type. When we consider a VOA $\mathbb{C}V$, we define the inner product on $\mathbb{C} \otimes_{\mathbb{R}} V_2$ by $\langle a, b \rangle \mathbf{1} = a_3 \bar{b}$, where $\bar{b}$ is a complex conjugate of b .

Under the above assumptions, $(V_2, \cdot(1)\cdot)$ becomes a commutative (non-associative) algebra called a Griess algebra with a positive definite invariant bilinear form. To simplify the notation, ef denotes $e(1)f$ for $e, f \in V_2$.

As we mentioned in the introduction, we add the following assumption.

(3) For any conformal vector e with central charge $\frac{1}{2}$, τ_e is not trivial.

In this case, there is a one-to-one correspondence between τ -involutions and conformal vectors with central charge $\frac{1}{2}$, see [Mi2].

Since V has a definite invariant bilinear form, if V contains a Virasoro VOA $L(c, 0)$, $L(c, 0)$ has a unitary highest weight representation in V . The work in [FQS] and [GKO] gives a complete classification of unitary highest weight representations of the Virasoro algebra. They proved that the highest weight representation $L(c, h)$ is unitary if and only if either (c, h) satisfies $c \geq 1$ and $h \geq 0$, or else (c, h) is among the following list:

$$\begin{aligned} c = c_m &= 1 - \frac{6}{(m+2)(m+3)} \quad (m = 0, 1, 2, \dots), \\ h = h_{r,s}^m &= \frac{[(m+3)r - (m+2)s]^2 - 1}{4(m+2)(m+3)} \quad (r, s \in \mathbb{N}, 1 \leq s \leq r \leq m+1). \end{aligned} \tag{2.1}$$

The unitary representations $L(c_m, h_{r,s}^m)$ for $(c_m, h_{r,s}^m)$ in the discrete series as above are called the discrete series of the Virasoro algebra.

Lemma 2.1 ([Mi2]) *V_2 decomposes into*

$$V_2 = \mathbb{R}e \oplus E^e(0) \oplus E^e\left(\frac{1}{2}\right) \oplus E^e\left(\frac{1}{16}\right),$$

where $E^e(h)$ denotes the eigenspace of $e(1)$ with eigenvalue h .

[Proof] As a $\text{VA}(e)$ -module, V decomposes into a direct sum of copies of $L(\frac{1}{2}, 0)$, $L(\frac{1}{2}, \frac{1}{2})$ and $L(\frac{1}{2}, \frac{1}{16})$ since $\text{VA}(e)$ is a rational VOA isomorphic to $L(\frac{1}{2}, 0)$ and $L(\frac{1}{2}, 0)$, $L(\frac{1}{2}, \frac{1}{2})$, $L(\frac{1}{2}, \frac{1}{16})$ are the set of irreducible modules. So V_2 decomposes into a direct sum of eigenspaces of $e(1)$ with eigenvalues $0, 2+r, \frac{1}{2}+r$ and $\frac{1}{16}+r$ ($r = 0, 1, 2, \dots$). If $e(1)v = (r+h)v$ for $v \in V_2$ and $r+h \neq 0, 2, \frac{1}{2}, \frac{1}{16}$, then v is not in the top module of $L(c, h)$ as a $\text{VA}(e)$ -module and so there is an element $u \in \text{VA}(e)_m$ such that $u(m)v \neq 0$ in $L(\frac{1}{2}, h)$, which contradicts $u_mv \in V_1 = \{0\}$. Since V has a definite invariant bilinear form, if a central charge of a conformal vector is less than one, it should be one of minimal discrete series. Therefore e is indecomposable and so $E^e(2) = \mathbb{R}e$, see [Mi2]. So we obtain the desired decomposition. ■

We note that from the fusion rules of $L(\frac{1}{2}, 0)$ -modules:

$$\begin{aligned}
L(\tfrac{1}{2}, h) \times L(\tfrac{1}{2}, k) &= L(\tfrac{1}{2}, k) \times L(\tfrac{1}{2}, h) \text{ for any } h, k = 0, \tfrac{1}{2}, \tfrac{1}{16} \\
L(\tfrac{1}{2}, 0) \times L(\tfrac{1}{2}, k) &= L(\tfrac{1}{2}, k) \text{ for any } k = 0, \tfrac{1}{2}, \tfrac{1}{16} \\
L(\tfrac{1}{2}, \tfrac{1}{2}) \times L(\tfrac{1}{2}, \tfrac{1}{2}) &= L(\tfrac{1}{2}, 0) \\
L(\tfrac{1}{2}, \tfrac{1}{2}) \times L(\tfrac{1}{2}, \tfrac{1}{16}) &= L(\tfrac{1}{2}, \tfrac{1}{16}) \quad \text{and} \\
L(\tfrac{1}{2}, \tfrac{1}{16}) \times L(\tfrac{1}{2}, \tfrac{1}{16}) &= L(\tfrac{1}{2}, 0) + L(\tfrac{1}{2}, \tfrac{1}{2})
\end{aligned}$$

we have

$$\begin{aligned}
ab &\in E^e(h) && \text{for } a \in \mathbb{R}e + E^e(0), b \in E^e(h) \text{ and any } h \\
ab &\in \mathbb{R}e + E^e(0) && \text{for } a, b \in E^e(\tfrac{1}{2}) \\
ab &\in E^e(\tfrac{1}{16}) && \text{for } a \in \mathbb{R}e + E^e(0) + E^e(\tfrac{1}{2}), b \in E^e(\tfrac{1}{16}) \text{ and} \\
ab &\in \mathbb{R}e + E^e(0) + E^e(\tfrac{1}{2}) && \text{for } a, b \in E^e(\tfrac{1}{16}).
\end{aligned}$$

Let e and f be two rational conformal vectors with central charge $\frac{1}{2}$ and assume that $\tau_e \tau_f$ is of order three. Set $G = \langle \tau_e, \tau_f \rangle$. G is isomorphic to a Symmetric group S_3 on three letters. The purpose in this section is to determine a subalgebra $\mathcal{G}$ of V_2 generated by e and f .

$(\tau_e \tau_f)^3 = \tau_e \tau_f \tau_e \tau_f \tau_e \tau_f = 1$ implies $\tau_e^{-1} \tau_f \tau_e = \tau_f^{-1} \tau_e \tau_f$. Since $\tau_{\phi(e)} = \phi \tau_e \phi^{-1}$ for any automorphism ϕ of V and τ gives arise a one-to-one correspondence between conformal vectors and τ -involutions by the assumption (3) and the statement after (3), we obtain

$$f^{\tau_e} = e^{\tau_f}.$$

We also note that $\tau_e^{-1} \tau_f \tau_e(e) = f$ and $\tau_e^{-1} \tau_f \tau_e(f) = e$. To simplify the notation, we assume that V coincides with a VOA $\text{VA}(\omega, e, f)$ generated by ω, e, f , where ω is the Virasoro element of V .

As it is well known, for a conformal vector v , its central charge is $2\langle v, v \rangle$. In particular, if e is a conformal vector with central charge $\frac{1}{2}$, then $\langle e, e \rangle = \frac{1}{4}$. Set $\langle e, f \rangle = \frac{\lambda}{4}$. Using the decomposition: $V_2 = \mathbb{R}e \oplus E^e(0) \oplus E^e(\frac{1}{2}) \oplus E^e(\frac{1}{16})$, we obtain

$$f = \lambda e + a + b + c$$

where $\lambda \in \mathbb{R}$, $a \in E^e(0)$, $b \in E^e(\frac{1}{2})$ and $c \in E^e(\frac{1}{16})$. By the definition of τ_e ,

$$f^{\tau_e} = \lambda e + a + b - c.$$

Similarly, we have

$$\begin{aligned}
e &= \lambda f + g + h + i && \text{and} \\
e^{\tau_f} &= \lambda f + g + h - i
\end{aligned}$$

where $g \in E^f(0)$, $h \in E^f(\frac{1}{2})$ and $i \in E^f(\frac{1}{16})$.

It follows from $e^{\tau f} = f^{\tau e}$ that $\lambda e + a + b - c = e - 2i$ and so we obtain

$$i = \frac{(1-\lambda)}{2}e - \frac{1}{2}a - \frac{1}{2}b + \frac{1}{2}c. \quad (2.1)$$

$ef = fe$ implies $2\lambda e + \frac{1}{2}b + \frac{1}{16}c = 2\lambda f + \frac{1}{2}h + \frac{1}{16}i$ and so we have

$$h = (4\lambda - \frac{1}{16})(1-\lambda)e + (\frac{1}{16} - 4\lambda)a + (\frac{17}{16} - 4\lambda)b + (\frac{1}{16} - 4\lambda)c \quad (2.2)$$

On the other hand, since $e = \lambda f + g + h + i$ and $f = \lambda e + a + b + c$, we obtain

$$g = (1-\lambda)(\frac{9}{16} - 3\lambda)e + (3\lambda + \frac{7}{16})a + (3\lambda - \frac{9}{16})b + (3\lambda - \frac{9}{16})c \quad (2.3)$$

By $ff = 2f$, we have

$$2\lambda e + 2a + 2b + 2c = 2\lambda^2 e + \lambda b + \frac{\lambda}{8}c + aa + bb + cc + 2ab + 2ac + 2bc.$$

Comparing the components in $E^e(\frac{1}{16})$, $\frac{\lambda}{8}c + 2(a+b)c - 2c = 0$ and so we get

$$(a+b)c = c - \frac{\lambda}{16}c. \quad (2.4)$$

Since $\langle e, aa \rangle = \langle ae, a \rangle = \langle 0, a \rangle = 0$, we have

$$aa \in E^e(0).$$

Since $bb \in \mathbb{R}e \oplus E^e(0)$, we denote it by

$$bb = (bb)_e e + (bb)_0,$$

where $(bb)_e \in \mathbb{C}$, $(bb)_0 \in E^e(0)$. Decompose cc as

$$cc = \{(2\lambda - 2\lambda^2 - (bb)_e)e\} + \{2a - aa - (bb)_0\} + \{(2 - \lambda - 2a)b\} \in \mathbb{R}e \oplus E^e(0) \oplus E^e(\frac{1}{2}). \quad (2.5)$$

From $fg = 0$,

$$0 = (\lambda e + a + b + c) \left((\lambda - 1)(3\lambda - \frac{9}{16})e + (3\lambda + \frac{7}{16})a + (3\lambda - \frac{9}{16})b + (3\lambda - \frac{9}{16})c \right).$$

The components in $E^e(\frac{1}{16})$ are

$$\begin{aligned} 0 &= \frac{\lambda}{16}(3\lambda - \frac{9}{16})c + (3\lambda - \frac{9}{16})(a+b)c + (\lambda - 1)(3\lambda - \frac{9}{16})\frac{1}{16}c \\ &\quad + (3\lambda + \frac{7}{16})(a+b)c - bc \end{aligned}$$

and so we have

$$bc = \frac{23}{28}(2^4\lambda - 1)c \quad (2.6)$$

and

$$ac = \frac{93}{2^8}(3 - 16\lambda)c. \quad (2.7)$$

The components in $E^e(\frac{1}{2})$ are

$$\begin{aligned} 0 = & \frac{\lambda}{2}(3\lambda - \frac{9}{16})b + (3\lambda - \frac{9}{16})ab + \frac{1}{2}(\lambda - 1)(3\lambda - \frac{9}{16})b \\ & + (3\lambda + \frac{7}{16})ab + (3\lambda - \frac{9}{16})(cc)_{\frac{1}{2}} \end{aligned}$$

where $(cc)_{\frac{1}{2}}$ denotes the component of cc in $E^e(\frac{1}{2})$, that is, $(2 - \lambda)b - 2ab$. Hence

$$\begin{aligned} 0 = & (\lambda(48\lambda - 9) + (96\lambda - 18)a + (48\lambda^2 - 71\lambda + 9) + (96\lambda + 14)a \\ & + (48\lambda - 9)(4 - 2\lambda - 4a))b \\ = & (48\lambda^2 - 9\lambda + 48\lambda^2 - 71\lambda + 9 + 192\lambda - 36 - 96\lambda^2 + 18\lambda \\ & + 192\lambda a - 4a - 192\lambda a + 36a) \\ = & (130\lambda - 27 + 32a)b \end{aligned}$$

and so we get

$$ab = \frac{9}{2^5}(3 - 2^4\lambda)b. \quad (2.8)$$

The components in $E^e(0)$ are

$$\begin{aligned} 0 = & (3\lambda + \frac{7}{16})aa + (3\lambda - \frac{9}{16})(bb)_0 + (3\lambda - \frac{9}{16})(cc)_0 \\ = & (3\lambda + \frac{7}{16})aa + (3\lambda - \frac{9}{16})(bb)_0 + (3\lambda - \frac{9}{16})(2a - aa - (bb)_0) \\ = & 3\lambda aa + \frac{7}{16}aa + 3\lambda(bb)_0 - \frac{9}{16}(bb)_0 + 3\lambda 2a - 3\lambda aa - 3\lambda(bb)_0 \\ & - 2a\frac{9}{16} + \frac{9}{16}aa + \frac{9}{16}(bb)_0 \\ = & aa + 6\lambda a - \frac{9}{8}a \end{aligned}$$

and so we obtain

$$aa = \frac{3}{2^3}(3 - 2^4\lambda)a. \quad (2.9)$$

Using $fh = \frac{1}{2}h$, we have

$$\begin{aligned} & (\lambda e + a + b + c)\{(\frac{65}{16}\lambda - 4\lambda^2 - \frac{1}{16})e + (\frac{1}{16} - 4\lambda)a \\ & + (\frac{17}{16} - 4\lambda)b + (\frac{1}{16} - 4\lambda)c\} \\ = & \frac{1}{2}\{(\frac{65}{16}\lambda - 4\lambda^2 - \frac{1}{16})e + (\frac{1}{16} - 4\lambda)a + (\frac{17}{16} - 4\lambda)b + (\frac{1}{16} - 4\lambda)c\}. \end{aligned}$$

Comparing the components in $W(0)$, we have

$$(\frac{1}{16} - 4\lambda)aa + (\frac{17}{16} - 4\lambda)(bb)_0 + (\frac{1}{16} - 4\lambda)(cc)_0 = (\frac{1}{32} - 2\lambda)a,$$

where $(cc)_0$ denotes the component of cc at $E^e(0)$. We hence obtain

$$(bb)_0 = \frac{3}{2^5}(2^6\lambda - 1)a. \quad (2.10)$$

Substituting these into the expansion of cc , we have

$$cc = (2\lambda - 2\lambda^2 - (bb)_e)e + \frac{31}{32}a + (8\lambda + \frac{5}{16})b \quad (2.11)$$

Therefore we obtain:

Lemma 2.2 $\mathcal{G} = \mathbb{R}e + \mathbb{R}a + \mathbb{R}b + \mathbb{R}c$ is a subalgebra. Furthermore, the symmetric group $G = \langle \tau_e, \tau_f \rangle$ acts on $\mathcal{G}$.

[**Proof**] We have already shown that $\mathcal{G}$ is a subalgebra. It is generated by e and f and it also contains e^{τ_f} . Hence $\mathcal{G}$ is invariant under the actions of τ_e and τ_f , that is, $\mathcal{G}$ is G -invariant. ■

3 Inner product

In this section, we will calculate the inner products of elements in $\mathcal{G}$ and show that λ is either $\frac{1}{2^6}$ or $\frac{13}{2^8}$.

Since

$$\langle ac, c \rangle = \langle a, cc \rangle = \frac{31}{32}\langle a, a \rangle$$

and

$$\langle ac, c \rangle = \frac{93}{2^8}(3 - 2^4\lambda)\langle c, c \rangle,$$

we have

$$\langle a, a \rangle = \frac{3}{8}(3 - 2^4\lambda)\langle c, c \rangle. \quad (3.1)$$

From

$$\langle bc, c \rangle = \langle b, cc \rangle = (8\lambda + \frac{5}{2^4})\langle b, b \rangle \text{ and } \langle bc, c \rangle = \frac{23}{2^8}(2^6\lambda - 1)\langle c, c \rangle,$$

we obtain

$$\langle b, b \rangle = \frac{23(2^6\lambda - 1)}{2^4(2^7\lambda + 5)}\langle c, c \rangle \quad (3.2)$$

We also have

$$\frac{9}{32}(3 - 2^4\lambda)\langle b, b \rangle = \langle ab, b \rangle = \langle a, bb \rangle = \frac{3}{2^5}(2^6\lambda - 1)\langle a, a \rangle. \quad (3.3)$$

Substituting (3.1) and (3.2) into (3.3),

$$(2^6\lambda - 1)(-2^4\lambda)\langle c, c \rangle = \frac{23(2^6\lambda - 1)}{2(2^7\lambda + 5)}\langle c, c \rangle. \quad (3.4)$$

If $\langle c, c \rangle = 0$, then $c = 0$ and so $f^{\tau_e} = f$ and $e = f^{\tau_e\tau_f} = f$, which contradicts $e \neq f$. Hence $\langle c, c \rangle \neq 0$ and λ is one of

$$\frac{3}{2^4}, \frac{1}{2^6}, \text{ or } \frac{13}{2^8}. \quad (3.5)$$

On the other hand, we obtain

$$\langle c, c \rangle = 16\langle ec, c \rangle = 16\langle e, cc \rangle = 4(2\lambda - 2\lambda^2 - (bb)_e). \quad (3.6)$$

Since

$$(\tau_e^{-1}\tau_f^{-1}\tau_e)\tau_e(\tau_e^{-1}\tau_f\tau_e) = \tau_e\tau_f\tau_e\tau_f\tau_e = \tau_f,$$

$\tau_e^{-1}\tau_f\tau_e(e) = f$ and $\tau_e^{-1}\tau_f\tau_e(f) = e$. Since c and i are uniquely defined by e and f , $\tau_e\tau_f\tau_e(c) = i$ and so we obtain $\langle c, c \rangle = \langle i, i \rangle$. Therefore,

$$\begin{aligned} \langle c, c \rangle &= \langle i, i \rangle = \frac{1}{4}\langle (1 - \lambda)e - a - b + c, (1 - \lambda)e - a - b + c \rangle \\ &= \frac{1}{4}\{(1 - \lambda)^2\frac{1}{4} + \langle a, a \rangle + \langle b, b \rangle + \langle c, c \rangle\} \\ &= \frac{1}{16}(1 - 2\lambda) + \frac{1}{16} = \frac{1}{8}(1 - \lambda) \end{aligned} \quad (3.7)$$

and so we have

$$\langle a, a \rangle = \frac{3(1 - \lambda)(3 - 2^4\lambda)}{2^6} \quad (3.8)$$

$$\langle b, b \rangle = \frac{23}{2^7} \frac{(1 - \lambda)(2^6\lambda - 1)}{2^7\lambda + 5} \quad \text{and} \quad (3.9)$$

$$\langle c, c \rangle = \frac{1 - \lambda}{8} \quad (3.10)$$

Using $\langle f, f \rangle = \frac{1}{4}$, we have

$$\begin{aligned} \frac{1}{4} &= \lambda^2\frac{1}{4} + \frac{-3 \cdot 2^4\lambda + 9}{8}\langle c, c \rangle + \frac{23(2^6\lambda - 1)}{(2^7\lambda + 5)2^4}\langle c, c \rangle + \langle c, c \rangle \\ &= \lambda^2\frac{1}{4} + \frac{-3 \cdot 2^4\lambda + 9}{8}\frac{1}{2}(1 - \lambda) + \frac{23(2^6\lambda - 1)}{(2^7\lambda + 5)2^4}\frac{1}{2}(1 - \lambda) + \frac{1}{2}(1 - \lambda) \end{aligned}$$

and

$$\begin{aligned}
0 &= (\lambda^2 - 1)\frac{1}{4} + \frac{-3 \cdot 2^4 \lambda + 9}{8} \frac{1}{2} (1 - \lambda) + \frac{23(2^6 \lambda - 1)}{(2^7 \lambda + 5)2^4} \frac{1}{2} (1 - \lambda) + \frac{1}{2} (1 - \lambda) \\
&= (1 - \lambda) \left\{ (-\lambda^2 - 1)\frac{1}{4} + \frac{-3 \cdot 2^4 \lambda + 9}{8} \frac{1}{2} + \frac{23(2^6 \lambda - 1)}{(2^7 \lambda + 5)2^4} \frac{1}{2} + \frac{1}{2} \right\} \\
&= \frac{(1 - \lambda)(2^6 \lambda - 1)(13 - 2^8 \lambda)}{2^4(2^7 \lambda + 5)}.
\end{aligned}$$

It follows from (3.5) that λ is either $\frac{1}{2^6}$ or $\frac{13}{2^8}$.

As we mentioned in the introduction, the both cases occur in the monstrous Griess algebra. So we have the following theorem.

Theorem 3.1 *Let V be a VOA of moonshine type satisfying the assumption (3) and let e and f be two distinct rational conformal vectors with central charge $\frac{1}{2}$. If $\tau_e \tau_f$ is of order three, then $\langle e, f \rangle$ is either $\frac{1}{2^8}$ or $\frac{13}{2^{10}}$.*

We next study a structure of a subVA $\text{VA}(e, f)$ generated by e and f . First we will show that $\text{VA}(e, f)$ has a Virasoro element and $\mathcal{G}$ contains it.

Since $aa = \frac{3}{8}(3 - 16\lambda)a$,

$$\omega_1 = \frac{16}{(9 - 48\lambda)}a \quad (3.11)$$

is a conformal vector.

Lemma 3.2 *$e + \omega_1$ is a Virasoro element of $\text{VA}(e, f)$.*

[Proof] By direct calculation using (2.7), (2.8), (2.9),

$$(e + \omega_1)e = 2e, \quad (e + \omega_1)a = 2a, \quad (e + \omega_1)b = 2b, \quad (e + \omega_1)c = 2c.$$

So it is sufficieint to show that $(e + \omega_1)(0)$ satisfies the derivation property for $v \in \text{VA}(e, f)$, that is, $Y((e + \omega_1)(0)v, z) = \frac{d}{dz}Y(v, z)$. As a $\text{VA}(e) \cong L(\frac{1}{2}, 0)$ -module, a is a highest weight vector in $L(\frac{1}{2}, 0)$ and so $e(0)a = 0$. By the skew-symmetry property, $a(0)e = 0$ and so

$$Y((e + \omega_1)(0)e, z) = Y(e(0)e, z) = \frac{d}{dz}Y(e, z).$$

Since $e + \omega_1$ is a unique element of $\mathcal{G}$ satisfying $(e + \omega_1)v = 2v$ for all $v \in \mathcal{G}$, G fixes it and so $Y((e + \omega_1)(0)f, z) = \frac{d}{dz}Y(f, z)$. It is shown in [Li] that if $a(z)$ and $b(z)$ satisfy D -derivation property, then so does $a(z)_n b(z)$ for any integer n , where $a(z)_n b(z)$ denotes n -th product. Hence $(e + \omega_1)(0)$ has the derivation property for all elements in $\text{VA}(e, f)$.

■

From now on, we assume that $V = \text{VA}(e, f)$ because it has a Virasoro element. Since it is convenient to use a linear representation of $\langle \tau_e \tau_f \rangle$, we will treat the complexification $\mathbb{C} \otimes_{\mathbb{R}} V$ of V . So from now on, V denotes $\mathbb{C} \otimes_{\mathbb{R}} V$.

4 The case $\lambda = \frac{1}{2^6}$

First we will study the case $\lambda = \frac{1}{2^6}$. In this case, $\langle b, b \rangle = 0$ and so $b = 0$. Namely, $\mathcal{G}$ is of dimension three. We will show $(\text{VA}(e, f))_2 = \mathcal{G}$.

The structure of $\mathcal{G}$ is given by

$$\begin{aligned} aa &= \frac{33}{32}a \\ ac &= \frac{7 \times 11 \times 13}{2^{10}}c \\ cc &= \frac{3^2 \times 7}{2^{11}}e + \frac{31}{32}a + \frac{7}{2^4}b. \end{aligned}$$

In $\mathcal{G} = \mathbb{C}e + \mathbb{C}a + \mathbb{C}c$, a Virasoro element decomposes into an orthogonal sum of two conformal vectors e with central charge $\frac{1}{2}$ and ω_1 with central charge $\frac{21}{22}$. The both $\text{VA}(e) \cong L(\frac{1}{2}, 0)$ and $\text{VA}(\omega_1) \cong L(\frac{21}{22}, 0)$ are minimal series and rational. In particular, ω_1 is indecomposable and so $E^e(0) = \mathbb{C}a$ by [Mi2].

Lemma 4.1 $E^e(\frac{1}{2}) = 0$.

[Proof] Viewing V as a $L(\frac{1}{2}, 0) \otimes L(\frac{21}{22}, 0)$ -module, V is a direct sum of irreducible $L(\frac{1}{2}, 0) \otimes L(\frac{21}{22}, 0)$ -modules. If $E^e(\frac{1}{2}) \neq 0$, then $L(\frac{21}{22}, 0)$ has to have an irreducible module with a highest weight $\frac{3}{2}$. However, there is no $\frac{3}{2}$ in the list of highest weights of $L(\frac{21}{22}, 0)$ -modules:

$$\frac{(12r - 11s)^2 - 1}{4 \times 11 \times 12} \quad 1 \leq s \leq r \leq 10.$$

■

Actually, the highest weights of irreducible $L(\frac{21}{22}, 0)$ -modules are

$$\begin{aligned} &0, \frac{3}{16 \times 11}, \frac{1}{16}, \frac{5}{2^4 \times 11}, \frac{1}{22}, \frac{35}{2^4 \times 3 \times 11}, \frac{1}{11}, \frac{21}{2^4 \times 11}, \frac{5}{33}, \frac{3}{16}, \frac{7}{22}, \\ &\frac{65}{2^4 \times 11}, \frac{14}{33}, \frac{85}{2^4 \times 11}, \frac{6}{11}, \frac{323}{2^4 \times 3 \times 11}, \frac{15}{22}, \frac{133}{2^4 \times 11}, \frac{5}{6}, \frac{13}{11}, \frac{225}{2^4 \times 11}, \frac{91}{2 \times 3 \times 11}, \\ &\frac{261}{2^4 \times 11}, \frac{35}{22}, \frac{899}{2^4 \times 3 \times 11}, \frac{20}{11}, \frac{31}{2^4}, \frac{57}{11}, \frac{481}{2^4 \times 11}, \frac{95}{33}, \frac{533}{2^4 \times 11}, \frac{35}{11}, \frac{1763}{2^4 \times 3 \times 11}, \\ &\frac{7}{2}, \frac{50}{11}, \frac{833}{2^4 \times 11}, \frac{325}{66}, \frac{901}{2^4 \times 11}, \frac{117}{22}, \frac{265}{2^4 \times 3}, \frac{155}{11}, \frac{1281}{2^4 \times 11}, \frac{248}{33}, \frac{1365}{2^4 \times 11}, \\ &8, \frac{121}{12}, \frac{1825}{2^4 \times 11}, \frac{703}{66}, \frac{175}{2^4}, \frac{301}{22}, \frac{2465}{2^4 \times 11}, \frac{43}{3}, \frac{196}{11}, \frac{291}{16}, \frac{45}{2} \end{aligned}$$

Since V has integer weights, V is a direct sum of copies of

$$\begin{aligned} &L(\frac{1}{2}, 0) \otimes L(\frac{21}{22}, 0), \quad L(\frac{1}{2}, 0) \otimes L(\frac{21}{22}, 8), \quad L(\frac{1}{2}, \frac{1}{2}) \otimes L(\frac{21}{22}, \frac{7}{2}), \\ &L(\frac{1}{2}, \frac{1}{2}) \otimes L(\frac{21}{22}, \frac{45}{2}), \quad L(\frac{1}{2}, \frac{1}{16}) \otimes L(\frac{21}{22}, \frac{31}{16}), \quad L(\frac{1}{2}, \frac{1}{16}) \otimes L(\frac{21}{22}, \frac{175}{16}) \end{aligned}$$

as a $L(\frac{1}{2}, 0) \otimes L(\frac{21}{22}, 0)$ -module.

Lemma 4.2 $V_2 = \mathcal{G}$ and $\dim V_2 = 3$.

[**Proof**] Suppose false and set $T = \mathcal{G}^\perp$. Since $\dim E^e(0) = 1$ and $E^e(\frac{1}{2}) = 0$, $T \subseteq E^e(\frac{1}{16})$ and so $ev = \frac{1}{16}v$ and $\tau_e(v) = -v$ for $v \in T$. Since G acts on T . $fv = (\frac{1}{26}e + a + c)v = \frac{1}{16}v$, which implies $cv = 0$ and $av = \frac{2^6-1}{2^{10}}$. Moreover, $\tau_e\tau_f$ acts T as 1. In particular, $vu \in (\mathbb{C}e + E^e(0))^G = \mathbb{C}\omega$ for $u, v \in T$. However, since $0 = \langle a, uu \rangle = \langle au, u \rangle = \frac{2^6-1}{2^{10}}\langle u, u \rangle$, we have a contradiction. ■

Theorem 4.3 Assume that $\tau_e\tau_f$ is of order three and $\langle e, f \rangle = \frac{1}{28}$. Then $\text{VA}(e, f)$ is a VOA with central charge $\frac{16}{11}$ and a Griess algebra $(\text{VA}(e, f))_2$ is of dimension 3. $\text{VA}(e, f)$ contains $L(\frac{1}{2}, 0) \otimes L(\frac{21}{22}, 0)$.

5 The case $\lambda = \frac{13}{2^8}$

We are very interested in this case because each edge in the diagram of $Y_{5,5,3}$ or the graph of 21 involutions is corresponding to this case.

As we showed in the previous section, the structure of $\mathcal{G}$ is given by

$$\begin{aligned} ab &= \frac{3^2 \times 5 \times 7}{2^9}b \\ aa &= \frac{3 \times 5 \times 7}{2^7}a \\ bb &= \frac{3^9}{2^{15}}e + \frac{3^3}{2^7}a \\ ac &= \frac{5 \times 7^2 \times 13}{2^{12}}c \\ bc &= \frac{3^2 \times 23}{2^{10}}c \\ cc &= \frac{3^5}{2^{13}}e + \frac{31}{32}a + \frac{23}{2^5}b. \end{aligned}$$

In particular,

$$\omega_1 = \frac{2^8}{105}a$$

is a conformal vector with central charge $\frac{81}{70}$.

Lemma 5.1 ω_1 is indecomposable. In particular, $E^e(0) = \mathbb{C}\omega_1$.

[Proof] Suppose that ω_1 is a sum $\omega' + \omega''$ of orthogonal conformal vectors. Since $c.c(\omega'), c.c(\omega'') \geq \frac{1}{2}$, $c.c(\omega'), c.c(\omega'') \leq \frac{23}{35}$, where $c.c(\omega')$ denotes a central charge of ω' . In particular, $\text{VA}(\omega')$ and $\text{VA}(\omega'')$ belong to minimal discrete series. However, the central charge of minimal discrete series which is less than or equal to $\frac{23}{35}$ are only $\{\frac{1}{2}, \frac{7}{10}\}$. However, $\frac{81}{70}$ is not a sum of them, which is a contradiction. ■

5.1 θ -fixed point space

In this subsection, we will study a θ -fixed point space and find conformal vectors of central charge $\frac{4}{5}$ and $\frac{6}{7}$. Using these conformal vectors, we will also show that $(\text{VA}(e, f))_2 = \mathcal{G}$.

Set $\theta = \tau_e \tau_f$, which is an automorphism of V of order three. We note that $e^\theta = f$ and $f^\theta = f^{\tau_e}$. Hence $\alpha = e + f + f^{\tau_e} = (\frac{13}{27} + 1)e + 2a + 2b$ and ω are in $\mathcal{G}^{<\theta>}$, where $\mathcal{G}^{<\theta>} = \{v \in \mathcal{G} | v^\theta = v\}$.

By the direct calculation, we have

$$\begin{aligned} ef &= \frac{13}{27}e + \frac{1}{2}b + \frac{1}{16}c \\ ff^{\tau_e} &= (\frac{13}{27} + \frac{13}{2^{12}} - \frac{1}{16})e + \frac{1}{16}a + \frac{9}{16}b \\ f^{\tau_e}e &= \frac{13}{27}e + \frac{1}{2}b - \frac{1}{16}c \end{aligned} \tag{5.1}$$

Hence

$$\alpha\alpha = 2e + 2f + 2f^{\tau_e} + (\frac{39}{26} + \frac{13}{2^{11}} - \frac{1}{8})e + \frac{1}{8}a + \frac{25}{16}b$$

and so

$$\alpha\alpha = \frac{57}{16}\alpha + (\frac{9 \times 13}{2^8} - \frac{27}{16})\omega. \tag{5.2}$$

Setting $\beta = \frac{24}{3}\alpha$, we obtain

$$\begin{aligned} \beta\beta &= 19\beta - 35\omega \\ \langle \beta, \beta \rangle &= \frac{47}{2} \\ \langle \beta, \omega \rangle &= 4 \quad \text{and} \\ \langle \omega, \omega \rangle &= \frac{29}{35}. \end{aligned}$$

It follows from a direct calculation that

$$\omega_2 = \frac{2(7\omega - \beta)}{9}$$

is a conformal vector with central charge $\frac{4}{5}$ and $\omega_3 = \omega - \omega_2$ is a conformal vector with central charge $\frac{6}{7}$. We note that the both belong to minimal discrete series. In particular, we have:

Proposition 5.2 *V contains a rational VOA $L(\frac{4}{5}, 0) \otimes L(\frac{6}{7}, 0)$ whose Virasoro element is equal to ω .*

We next view V as a $L(\frac{4}{5}, 0) \otimes L(\frac{6}{7}, 0)$ -module. Let ϵ be a cubic root of unity and set $\gamma = e + \epsilon f + \epsilon^2 f^{\tau_e}$. Namely, $\gamma^\theta = \epsilon^{-1} \gamma$. Then we have

$$\begin{aligned} \alpha\gamma &= (e + f + f^{\tau_e})(e + \epsilon f + \epsilon^2 f^{\tau_e}) \\ &= 2(e + \epsilon f + \epsilon^2 f^{\tau_e}) + (\epsilon + 1)ef + (\epsilon^2 + 1)ef^{\tau_e} + (\epsilon + \epsilon^2)f f^{\tau_e} \\ &= 2\beta + \frac{1}{16}\gamma = \frac{33}{16}\gamma, \end{aligned}$$

which implies

$$\beta\gamma = \left(\frac{2^4}{3}\alpha\right)\gamma = 11\gamma$$

and

$$\omega_2\gamma = \frac{2}{3}\gamma.$$

We note $\{e, f, f^{\tau_e}\}$ is a linearly independent set and so $\gamma \neq 0$. $\theta = \tau_e \tau_f$ acts on $\mathbb{C}\gamma$ as ϵ^{-1} . Similarly, $\omega_2\gamma = \frac{2}{3}\gamma$. Therefore, V contains $U^1 \oplus U^2$ where $U^i \cong L(\frac{4}{5}, \frac{2}{3}) \otimes L(\frac{6}{7}, \frac{4}{3})$ and $\theta = \tau_e \tau_f$ acts on U^i as ϵ^i and τ_e exchange U^1 and U^2 .

Therefore,

$$e, f \in L(\frac{4}{5}, 0) \otimes L(\frac{6}{7}, 0) + L(\frac{4}{5}, \frac{2}{3}) \otimes L(\frac{6}{7}, \frac{4}{3}) + L(\frac{4}{5}, \frac{2}{3}) \otimes L(\frac{6}{7}, \frac{4}{3}).$$

From fusion rules of $L(\frac{4}{5}, 0)$ -modules:

$$\begin{aligned} L(\frac{4}{5}, \frac{2}{3}) \times L(\frac{4}{5}, \frac{2}{3}) &= L(\frac{4}{5}, 0) + L(\frac{4}{5}, \frac{2}{3}) + L(\frac{4}{5}, 3) \\ L(\frac{4}{5}, 3) \times L(\frac{4}{5}, \frac{2}{3}) &= L(\frac{4}{5}, \frac{2}{3}) \\ L(\frac{4}{5}, 3) \times L(\frac{4}{5}, 3) &= L(\frac{4}{5}, 0), \end{aligned}$$

we obtain that $V = \text{VA}(e, f)$ is a direct sum of copies of $L(\frac{4}{5}, 0)$, $L(\frac{4}{5}, \frac{2}{3})$ and $L(\frac{4}{5}, 3)$ as a $L(\frac{4}{5}, 0)$ -module, see Appendix.

The set of highest weights of irreducible $L(\frac{6}{7}, 0)$ -modules are

$$\frac{(7r - 6s)^2 - 1}{4 \times 6 \times 7} \quad 1 \leq s \leq r \leq 5,$$

that is,

$$\frac{1}{56}, \frac{1}{21}, \frac{5}{56}, \frac{1}{7}, \frac{3}{8}, \frac{10}{21}, \frac{33}{56}, \frac{5}{7}, \frac{4}{3}, \frac{85}{56}, \frac{12}{7}, \frac{23}{8}, \frac{22}{7}, \text{ and } 5.$$

Since the weights in V are integers, the possibilities of $L(\frac{4}{5}, 0) \otimes L(\frac{6}{7}, 0)$ -submodules are

$$\begin{aligned} &L(\frac{4}{5}, 0) \otimes L(\frac{6}{7}, 0), \quad L(\frac{4}{5}, 0) \otimes L(\frac{6}{7}, 5), \\ &L(\frac{4}{5}, \frac{2}{3}) \otimes L(\frac{6}{7}, \frac{4}{3}), \\ &L(\frac{4}{5}, 3) \otimes L(\frac{6}{7}, 0), \quad L(\frac{4}{5}, 3) \otimes L(\frac{6}{7}, 5). \end{aligned}$$

The fusion rules for $L(\frac{6}{7}, 0)$ -modules $L(\frac{6}{7}, \frac{4}{3})$ and $L(\frac{6}{7}, 5)$ are

$$\begin{aligned} &L(\frac{6}{7}, 5) \times L(\frac{6}{7}, 5) = L(\frac{6}{7}, 0) \\ &L(\frac{6}{7}, 5) \times L(\frac{6}{7}, \frac{4}{3}) = L(\frac{6}{7}, \frac{4}{3}) \\ &L(\frac{6}{7}, \frac{4}{3}) \times L(\frac{6}{7}, \frac{4}{3}) = L(\frac{6}{7}, 0) + L(\frac{6}{7}, \frac{4}{3}) + L(\frac{6}{7}, 5), \end{aligned}$$

see [W].

Our last purposes is to show the following theorem.

Theorem 5.3 $(\text{VA}(e, f))_2 = \mathcal{G}$.

[**Proof**] Assume $\mathcal{G} \neq V_2$ and set $T = \mathcal{G}^\perp$. First we will show the following lemma.

Lemma 5.4 $T^{<\theta>} = 0$ and $(V_2)^{<\theta>} = \mathbb{C}\omega_2 + \mathbb{C}\omega_3$.

[**Proof**] Clearly, V contains only one copy of $L(\frac{4}{5}, 0) \otimes L(\frac{6}{7}, 0)$. So the eigen value of ω_2 on T is only $\frac{2}{3}$. It is clear that G acts on T . Suppose $P(\frac{1}{2}) = T^\theta \cap E^e(\frac{1}{2}) \neq 0$. Namely, $\tau_e, \tau_f, \tau_{f\tau_e}$ act $P(\frac{1}{2})$ as 1. In this case, for $u \in P(\frac{1}{2})$, $eu = \frac{1}{2}u$, $fu = \frac{1}{2}u$, $f^{\tau_e}u = \frac{1}{2}u$. Hence $cu = 0$. Moreover, $bu \in (\mathbb{C}e + E^e(0)) \cap T = \{0\}$. Since

$$2u = \omega u = (e + \omega_1)u = (e + \frac{2^8 a}{105})u = \frac{1}{2}u + \frac{2^8}{105}au,$$

we obtain $au = \frac{315}{2^9}u$. However, we have

$$\frac{1}{2}u = fu = (\frac{13}{2^8}e + a + b + c)u = \frac{13}{2^9}u + \frac{315}{2^9}u = \frac{328}{512}u,$$

a contradiction. We hence have $P(\frac{1}{2}) = 0$. We next assume $P(\frac{1}{16}) = T^\theta \cap E^e(\frac{1}{16}) \neq 0$. Namely, $\tau_e, \tau_f, \tau_{f\tau_e}$ act on $P(\frac{1}{16})$ as -1 . In this case, for $u \in P(\frac{1}{16})$, $eu = \frac{1}{16}u$, $fu = \frac{1}{16}u$, $f^{\tau_e}u = \frac{1}{16}u$ by the definition of τ -automorphism. Hence $cu = \frac{1}{2}(f - f^{\tau_e})u = 0$ and $\alpha u = \{(2\lambda + 1)e + 2a + 2b\}u = eu + 2(\lambda e + a + b)u = \frac{3}{16}u$. Since $\omega_2 = \frac{2(21\omega - 16\alpha)}{27}$,

we have $\frac{2}{3}u = \omega_2 u = \frac{2(\beta - 7\omega)}{-9}u = \frac{2(1 - 14)}{-9}u$, a contradiction. Therefore $T^\theta = 0$ and it is clear that $\mathcal{G}^{<\theta>} = \mathbb{C}\omega_2 + \mathbb{C}\omega_3$. ■

We next view V as a G -module. The symmetric group G on three letters has 3 irreducible modules $\mathbb{C}(+)$, $\mathbb{C}(-)$ and D , where $\mathbb{C}(+)$ is a trivial module, θ and τ_e act on $\mathbb{C}(-)$ as 1 and -1 , respectively, and D is an irreducible module of dimension two. Therefore, V has a subVOA V^G such that V decomposes into the direct sum

$$V = V^G \oplus U \otimes \mathbb{C}(-) \oplus W \otimes D,$$

where U and W are irreducible V^G -modules, see [DM].

Since $\omega_2, \omega_3 \in \text{VA}(e, f)^G$, V^G contains $L(\frac{4}{5}, 0) \otimes L(\frac{6}{7}, 0)$.

Recall $\gamma = e + \epsilon f + \epsilon^2 f^{\tau_e}$ and $\theta = \tau_e \tau_f$ acts γ as ϵ^{-1} . It is easy to check that $\gamma \neq 0$. Therefore W contains $L(\frac{4}{5}, \frac{2}{3}) \otimes L(\frac{6}{7}, \frac{4}{3})$. Lemma 5.3 implies the highest weight of U is greater than 3.

Case 1.

First assume $V^G = L(\frac{4}{5}, 0) \otimes L(\frac{6}{7}, 0)$. Then $W = L(\frac{4}{5}, \frac{2}{3}) \otimes L(\frac{6}{7}, \frac{4}{3})$ and so $V_2 = (L(\frac{4}{5}, 0) \otimes L(\frac{6}{7}, 0))_2 \oplus (W \otimes D)_2$ is of dimension four. Namely we have the desired result.

Case 2.

Assume $V^G \neq L(\frac{4}{5}, 0) \otimes L(\frac{6}{7}, 0)$. Then

$$V^{<\theta>} = (\frac{4}{5}, 0) \otimes L(\frac{6}{7}, 0) \oplus L(\frac{4}{5}, 0) \otimes L(\frac{6}{7}, 5) \oplus L(\frac{4}{5}, 3) \otimes L(\frac{6}{7}, 0) \oplus L(\frac{4}{5}, 3) \otimes L(\frac{6}{7}, 5),$$

which is a subVOA. In particular, $L(\frac{6}{7}, 0) \oplus L(\frac{6}{7}, 5)$ has a VOA structure.

$L(\frac{4}{5}, 0) \oplus L(\frac{4}{5}, 3)$ is a rational VOA and its irreducible modules are classified in [KMiy]. Namely, it has exactly the following six irreducible modules:

$$W(0), W(\frac{2}{5}), W(\frac{2}{3}, +), W(\frac{1}{15}, +), W(\frac{2}{3}, -), W(\frac{1}{15}, -).$$

Here h in $W(h)$ and $W(h, \pm)$ denotes the lowest degree and $W(k, -)$ is the contragredient (dual) module of $W(k, +)$ for $k = \frac{2}{3}, \frac{1}{15}$. As $L(\frac{4}{5}, 0)$ -modules,

$$\begin{aligned} W(0) &\cong L(\frac{4}{5}, 0) \oplus L(\frac{4}{5}, 3), \\ W(\frac{2}{5}) &\cong L(\frac{4}{5}, \frac{2}{5}) \oplus L(\frac{4}{5}, \frac{7}{5}), \\ W(\frac{2}{3}, +) &\cong L(\frac{4}{5}, \frac{2}{3}), \\ W(\frac{2}{3}, -) &\cong L(\frac{4}{5}, \frac{2}{3}), \\ W(\frac{1}{15}, +) &\cong L(\frac{4}{5}, \frac{1}{15}), \\ W(\frac{1}{15}, -) &\cong L(\frac{4}{5}, \frac{1}{15}). \end{aligned}$$

For an irreducible $L(\frac{6}{7}, 0) \oplus L(\frac{6}{7}, 5)$ -module, we will need the following lemma.

Lemma 5.5 *If $L(\frac{6}{7}, 0) \oplus L(\frac{6}{7}, 5)$ is a VOA and X is an irreducible module contains $L(\frac{6}{7}, \frac{4}{3})$, then $X \cong L(\frac{6}{7}, \frac{4}{3})$ as $L(\frac{6}{7}, 0)$ -modules.*

[Proof] Suppose false and let Y^X be a vertex operator of $L(\frac{6}{7}, 0) \oplus L(\frac{6}{7}, 5)$ on X . Since $L(\frac{6}{7}, 0)$ is rational, X is a direct sum of $L(\frac{6}{7}, 0)$ -modules. By the fusion rule $L(\frac{6}{7}, 5) \times L(\frac{6}{7}, \frac{4}{3}) = L(\frac{6}{7}, \frac{4}{3})$, we have

$$X = L(\frac{6}{7}, \frac{4}{3}) \oplus L(\frac{6}{7}, \frac{4}{3}).$$

Choose $0 \neq I \in I(\begin{smallmatrix} L(\frac{6}{7}, \frac{4}{3}) \\ L(\frac{6}{7}, 5) \end{smallmatrix} \begin{smallmatrix} L(\frac{6}{7}, \frac{4}{3}) \\ L(\frac{6}{7}, \frac{4}{3}) \end{smallmatrix})$. Since $\dim I(\begin{smallmatrix} L(\frac{6}{7}, \frac{4}{3}) \\ L(\frac{6}{7}, 5) \end{smallmatrix} \begin{smallmatrix} L(\frac{6}{7}, \frac{4}{3}) \\ L(\frac{6}{7}, \frac{4}{3}) \end{smallmatrix}) = 1$, there is $\mu \in \mathbb{C}$ such that $Y^X(u, z) = \begin{pmatrix} 0 & I(u, z) \\ \mu I(u, z) & 0 \end{pmatrix}$ for $u \in L(\frac{6}{7}, 5)$ by choosing a suitable basis. Replacing I by $\sqrt{\mu}I$, we have $Y^X(u, z) = \begin{pmatrix} 0 & I(u, z) \\ I(u, z) & 0 \end{pmatrix}$. Then $\{(w, w) \in L(\frac{6}{7}, \frac{4}{3}) \oplus L(\frac{6}{7}, \frac{4}{3})\}$ is a $L(\frac{6}{7}, 0) \oplus L(\frac{6}{7}, 5)$ -submodule, which contradicts the assumption on X . ■

Now we go back to the proof of Theorem 5.3. Viewing V as a $\langle \theta \rangle$ -module, $V = V^0 \oplus V^1 \oplus V^2$, where θ acts on V^i as $(\epsilon)^i$. Since

$$V^0 \cong (L(\frac{4}{5}, 0) \oplus L(\frac{4}{5}, 3)) \otimes (L(\frac{6}{7}, 0) \oplus L(\frac{6}{7}, 5)),$$

we have

$$V^i \cong L(\frac{4}{5}, \frac{2}{3}) \otimes L(\frac{6}{7}, \frac{4}{3})$$

and so $\dim V_2 = 4$ and $V_2 = \mathcal{G}$. This completes the proof of Theorem 5.3. ■

On $X = L(\frac{6}{7}, \frac{4}{3})$, there are two $L(\frac{6}{7}, 0) \oplus L(\frac{6}{7}, 5)$ -module structures. Namely, if $Y^X(v, z)$ is a vertex operator of $v \in L(\frac{6}{7}, 0) \oplus L(\frac{6}{7}, 5)$, then the other is defined by $Y^X(v, z)$ for $v \in L(\frac{6}{7}, 0)$ and $-Y^X(v, z)$ for $v \in L(\frac{6}{7}, 5)$. We denote them by $L(\frac{6}{7}, \frac{4}{3})^{\pm 1}$.

Theorem 5.6 *If $|\tau_e \tau_f| = 3$ and $\langle e, f \rangle = \frac{13}{2^{10}}$, then $\text{VA}(e, f)$ is isomorphic to one of following:*

- (1) $(0, 0) \oplus (3, 0) \oplus (W(\frac{2}{3}, +) \otimes L(\frac{6}{7}, \frac{4}{3})) \oplus (W(\frac{2}{3}, -) \otimes L(\frac{6}{7}, \frac{4}{3}))$
- (2) $(0, 0) \oplus (0, 5) \oplus (L(\frac{4}{5}, \frac{2}{3}) \otimes L(\frac{6}{7}, \frac{4}{3})^{+1}) \oplus (L(\frac{4}{5}, \frac{2}{3}) \otimes L(\frac{6}{7}, \frac{4}{3})^{-1})$
- (3) $(0, 0) \oplus (3, 5) \oplus (W(\frac{2}{3}, +) \otimes L(\frac{6}{7}, \frac{4}{3})^{+1}) \oplus (W(\frac{2}{3}, -) \otimes L(\frac{6}{7}, \frac{4}{3})^{+1})$
- (4) $(0, 0) \oplus (3, 0) \oplus (0, 5) \oplus (3, 5) \oplus (W(\frac{2}{3}, \pm) \otimes L(\frac{6}{7}, \frac{4}{3})^{+1}) \oplus (W(\frac{2}{3}, \mp) \otimes L(\frac{6}{7}, \frac{4}{3})^{-1})$
in (4), $\tau_e \tau_f$ is equal to an automorphism determined by a 3-State Potts model
 $L(\frac{4}{5}, 0) \oplus L(\frac{4}{5}, 3)$ defined in [Mi3]

where (h, k) denotes $L(\frac{4}{5}, h) \otimes L(\frac{6}{7}, k)$. We will call them VOAs of involution type A_2 .

At last we expect the following conjecture.

Conjecture 1 *Let e and f be two distinct rational conformal vectors with central charge $\frac{1}{2}$. If $\langle e, f \rangle = \frac{13}{28}$, then $\tau_e \tau_f$ is of order three.*

6 Appendix

Fusion rule of $L(\frac{4}{5}, 0)$ -modules

0	$\frac{2}{5}$	$\frac{1}{40}$	$\frac{7}{5}$	$\frac{21}{40}$	$\frac{1}{15}$	3	$\frac{13}{8}$	$\frac{2}{3}$	$\frac{1}{8}$
$\frac{2}{5}$	$0 : \frac{7}{5}$	$\frac{1}{8} : \frac{21}{40}$	$\frac{2}{5} : 3$	$\frac{1}{40} : \frac{13}{8}$	$\frac{1}{15} : \frac{2}{3}$	$\frac{7}{5}$	$\frac{21}{40}$	$\frac{1}{15}$	$\frac{1}{40}$
$\frac{1}{40}$	$\frac{1}{8} : \frac{21}{40}$	$0 : \frac{7}{5} : \frac{2}{3} : \frac{1}{15}$	$\frac{1}{40} : \frac{13}{8}$	$\frac{2}{5} : 3 : \frac{1}{15} : \frac{2}{3}$	$\frac{1}{40} : \frac{13}{8} : \frac{21}{40} : \frac{1}{8}$	$\frac{21}{40}$	$\frac{7}{5} : \frac{1}{15}$	$\frac{21}{40} : \frac{1}{40}$	$\frac{1}{15} : \frac{2}{5}$
$\frac{7}{5}$	$\frac{2}{5} : 3$	$\frac{1}{40} : \frac{13}{8}$	$0 : \frac{7}{5}$	$\frac{1}{8} : \frac{21}{40}$	$\frac{2}{3} : \frac{1}{15}$	$\frac{2}{5}$	$\frac{1}{40}$	$\frac{1}{15}$	$\frac{21}{40}$
$\frac{21}{40}$	$\frac{1}{40} : \frac{13}{8}$	$\frac{2}{5} : 3 : \frac{1}{15} : \frac{2}{3}$	$\frac{1}{8} : \frac{21}{40}$	$0 : \frac{7}{5} : \frac{2}{3} : \frac{1}{15}$	$\frac{1}{8} : \frac{21}{40} : \frac{13}{8} : \frac{1}{40}$	$\frac{1}{40}$	$\frac{2}{5} : \frac{1}{15}$	$\frac{1}{40} : \frac{21}{40}$	$\frac{1}{15} : \frac{7}{5}$
$\frac{1}{15}$	$\frac{1}{15} : \frac{2}{3}$	$\frac{1}{40} : \frac{13}{8} : \frac{21}{40} : \frac{1}{8}$	$\frac{2}{3} : \frac{1}{15}$	$\frac{1}{8} : \frac{21}{40} : \frac{13}{8} : \frac{1}{40}$	$0 : \frac{7}{5} : \frac{2}{3} : \frac{1}{15} : 3 : \frac{2}{5}$	$\frac{1}{15}$	$\frac{1}{40} : \frac{21}{40}$	$\frac{2}{5} : \frac{1}{15} : \frac{7}{5}$	$\frac{1}{40} : \frac{21}{40}$
3	$\frac{7}{5}$	$\frac{21}{40}$	$\frac{2}{5}$	$\frac{1}{40}$	$\frac{1}{15}$	0	$\frac{1}{8}$	$\frac{2}{3}$	$\frac{13}{8}$
$\frac{13}{8}$	$\frac{21}{40}$	$\frac{7}{5} : \frac{1}{15}$	$\frac{1}{40}$	$\frac{2}{5} : \frac{1}{15}$	$\frac{1}{40} : \frac{21}{40}$	$\frac{1}{8}$	$0 : \frac{2}{3}$	$\frac{1}{8} : \frac{13}{8}$	$\frac{2}{3} : 3$
$\frac{2}{3}$	$\frac{1}{15}$	$\frac{21}{40} : \frac{1}{40}$	$\frac{1}{15}$	$\frac{1}{40} : \frac{21}{40}$	$\frac{2}{5} : \frac{1}{15} : \frac{7}{5}$	$\frac{2}{3}$	$\frac{1}{8} : \frac{13}{8}$	$0 : \frac{2}{3} : 3$	$\frac{1}{8} : \frac{13}{8}$
$\frac{1}{8}$	$\frac{1}{40}$	$\frac{1}{15} : \frac{2}{5}$	$\frac{21}{40}$	$\frac{1}{15} : \frac{7}{5}$	$\frac{1}{40} : \frac{21}{40}$	$\frac{13}{8}$	$\frac{2}{3} : 3$	$\frac{1}{8} : \frac{13}{8}$	$0 : \frac{2}{3}$

Fusion rules of $L(\frac{4}{5}, 0) \oplus L(\frac{4}{5}, 3)$ -modules.

$W(0)$	$W(\frac{2}{5})$	$W(\frac{2}{3}, +)$	$W(\frac{1}{15}, +)$	$W(\frac{2}{3}, -)$	$W(\frac{1}{15}, -)$
$W(\frac{2}{5})$	$W(0) : W(\frac{2}{5})$	$W(\frac{1}{15}, +)$	$W(\frac{1}{15}, +) : W(\frac{2}{3}, +)$	$W(\frac{1}{15}, -)$	$W(\frac{1}{15}, -) : W(\frac{2}{3}, -)$
$W(\frac{2}{3}, +)$	$W(\frac{1}{15}, +)$	$W(\frac{2}{3}, -)$	$W(\frac{1}{15}, -)$	$W(0)$	$W(\frac{2}{5})$
$W(\frac{1}{15}, +)$	$W(\frac{1}{15}, +) : W(\frac{2}{3}, +)$	$W(\frac{1}{15}, -)$	$W(\frac{1}{15}, -) : W(\frac{2}{3}, -)$	$W(\frac{2}{5})$	$W(0) : W(\frac{2}{5})$
$W(\frac{2}{3}, -)$	$W(\frac{1}{15}, -)$	$W(0)$	$W(\frac{2}{5})$	$W(\frac{2}{3}, +)$	$W(\frac{1}{15}, +)$
$W(\frac{1}{15}, -)$	$W(\frac{1}{15}, -) : W(\frac{2}{3}, -)$	$W(\frac{2}{5})$	$W(0) : W(\frac{2}{5})$	$W(\frac{1}{15}, +)$	$W(\frac{1}{15}, +) : W(\frac{2}{3}, +)$

References

- [At] J. H. Conway, R. T. Curtis, S. P. Norton, R. A. Parker and R. A. Wilson, *ATLAS of Finite Groups*, Oxford Univ. Press, 1985.
- [Bo] R. E. Borcherds, Vertex algebra, Kac-Moody algebra, and the Monster, *Proc. Natl. Acad. Sci. USA* **83** (1986), 3068-3071.
- [Co] J. H. Conway, A simple construction for the Fischer-Griess monster group, *Invent. Math.* **79** (1985), 513-540.
- [CN] J. H. Conway and S. P. Norton, Monstrous moonshine, *Bull. London Math. Soc.* **11** (1979) 308-339.
- [CS] J. H. Conway and N. J. A. Sloane, *Sphere Packings, Lattices and Groups*, Springer-Verlag, New-York, (1988).

- [DM] C. Dong and G. Mason, On quantum Galois theory, *Duke Math. J.* **86** (1997), no.2, 305-321.
- [F] B.Fischer, Finite groups generated by 3-transpositions, *Invent. Math.* **13** (1971), 232-246
- [FLM] I. Frenkel, J. Lepowsky and A. Meurman, Vertex Operator Algebras and the Monster, *Pure and Appl. Math.* **Vol. 134**, Academic Press, Boston, 1988.
- [FQS] D. Friedan, Z.Qiu and S. Shenker, Conformal invariance, unitarity and two-dimensional critical exponents, *MSRI publ. # 3*, Springer-Verlag, (1984), 419-449.
- [GKO] P.Goddard, A. Kent and D. Olive, Unitary representations of the Virasoro algebra and super-Virasoro algebras, *Commun. Math. Phys.* **103** (1986), 105-119.
- [Gr] R. L. Griess, Jr., The friendly giant, *Invent. Math.* **69** (1982), 1-102.
- [GH] R. L. Griess, Jr. and G. Höhn, Virasoro frames and their stabilizers for the E_8 lattice type vertex operator algebra, *Crelle's Journal*, to appear.
- [KM] M. Kitazume and M. Miyamoto, 3-transposition automorphism group of VOAs, Groups and Combinatorics, *Advanced Studies in Pure Math.* **32**, (2001), 315-324.
- [KMiy] M. Kitazume, M. Miyamoto and H. Yamada, Ternary codes and vertex operator algebras, *J. Algebra*, **223** (2000), no. 2, 379-395.
- [Li] H.-S. Li, Local systems of vertex operators, vertex superalgebras and modules, *J. Pure Appl. Alg.*, **109** (1996), no.2, 143-195
- [Mi1] M. Miyamoto, 21 involutions acting on the Moonshine module, *J. Algebra* **175**, (1995), no. 3, 941-965.
- [Mi2] M. Miyamoto, Griess algebras and conformal vectors in vertex operator algebras, *J. Algebra* **179**, (1996) 523-548
- [Mi3] M. Miyamoto, 3-state Potts model and automorphisms of vertex operator algebras of order 3, *J. Algebra* **239** (2001), no. 1, 56-76.
- [MN] W. Meyer and W. Neutsch, Associative subalgebras of the Griess algebra, *J. of Algebra* **158** (1993), 1-17.
- [W] W. Wang, Rationality of Virasoro vertex operator algebra, *Duke Math. J. IMRN*, **Vol. 71**, No. 1 (1993), 197-211.